


# Does the 220 PeV Event at KM3NeT Point to New Physics?


Vedran Brdar  
Oklahoma State University





# Neutrino Astronomy: IceCube



# Neutrino Astronomy: IceCube




IceCube collaboration, Science 2023




# Former Energy Champion: Glashow Resonance at IceCube

Bull. Am. Astron. Soc. 51, 185 (2019)




- ▶ shower with an energy of  $6.05 \pm 0.72$  PeV
- ▶ presence of electron antineutrinos in the astrophysical flux  $\rightarrow$  way to distinguish  $\nu$  from  $\bar{\nu}$

- ▶  $\mathcal{O}(100)$  upgoing tracks and HESE events
- ▶ Glashow resonance: cross section enhancement from on-shell  $W^-$  production
- ▶  $\sigma \propto \frac{1}{(E-E_0)^2 + \Gamma^2}$ , with  $E_0 = \frac{M_W^2}{2m_e} \approx 6.3$  PeV





Nature 591, 220–224 (2021)






# KM3NeT





# New Energy Champion: KM3-230213A Event

- ▶ The KM3NeT collaboration reported the detection of a  $\sim 120$  PeV muon originating from a neutrino interaction with a **median neutrino energy of 220 PeV**
- ▶ This is the **highest-energy neutrino** ever detected, exceeding the Glashow resonance event in IceCube's dataset by a factor of  $\mathcal{O}(10)$



# What about IceCube?


- ▶ IceCube has been operating with a much larger effective area for a longer time and **has not observed neutrinos above  $\sim 10$  PeV**
- ▶  **$2\text{--}3.5\sigma$  tension**, depending on the neutrino source (Li et al., arXiv:2502.04508)
- ▶ Event such as KM3-230213A would be expected in 70 years of observation...an upward fluctuation at the level of  $2.2\sigma$  (KM3NeT, Nature (2025))




# KM3NeT vs IceCube

$$\frac{dN(E_\nu)}{dE} = T \int d\Omega A_{\text{eff}}(E_\nu, \cos\theta) \Phi(E_\nu, \Omega)$$

- ▶ The difference in the effective areas between IceCube and KM3NeT is  $\sim 20$
- ▶ To explain the tension, the neutrino flux at KM3NeT **needs to be larger by a similar factor**
- ▶ How to achieve that?  
 $\implies$  **Sterile neutrinos** partially converting into active neutrinos inside Earth





# Sterile Neutrino Sources



- ▶ Lack of multi-messenger observations corresponding to KM3-230213A event

Speculations on sterile neutrino sources:

- ▶ A dense outer layer around the source - stops/downscatters SM particles
- ▶ AGN jets + dense matter blocking ultra high energy SM particles
- ▶ Dark sector stars



arXiv > hep-ph > arXiv:2502.21299

Search...

Help


High Energy Physics - Phenomenology

[Submitted on 28 Feb 2025]

## Does the 220 PeV Event at KM3NeT Point to New Physics?

Vedran Brdar, Dibya S. Chattopadhyay

The KM3NeT collaboration recently reported the observation of KM3-230213A, a neutrino event with an energy exceeding 100 PeV, more than an order of magnitude higher than the most energetic neutrino in IceCube's catalog. Given its longer data-taking period and larger effective area relative to KM3NeT, IceCube should have observed events around that energy. This tension has recently been quantified to lie between  $2\sigma$  and  $3.5\sigma$ , depending on the neutrino source. A  $\mathcal{O}(100)$  PeV neutrino detected at KM3NeT has traversed approximately 147 km of rock and sea en route to the detector, whereas neutrinos arriving from the same location in the sky would have only traveled through about 14 km of ice before reaching IceCube. We use this difference in propagation distance to address the tension between KM3NeT and IceCube. Specifically, we consider a scenario in which the source emits sterile neutrinos that partially convert to active neutrinos through oscillations. We scrutinize two such realizations, one where a new physics matter potential induces a resonance in sterile-to-active transitions and another one where off-diagonal neutrino non-standard interactions are employed. In both cases, sterile-to-active neutrino oscillations become relevant at length scales of  $\sim 100$  km, resulting in increased active neutrino flux near the KM3NeT detector, alleviating the tension between KM3NeT and IceCube. Overall, we propose the exciting possibility that neutrino telescopes may have started detecting new physics.



- ▶  $\mathcal{O}(100)$  PeV neutrino detected at KM3NeT has traversed  $\sim 150$  km of rock and sea en route to the detector
- ▶ Neutrinos arriving from the same location in the sky (angle of  $8^\circ$  w.r.t. horizon) would have only traveled through  $\sim 14$  km of ice before reaching IceCube

# Model 1



# Neutrino production through matter-induced resonance

- ▶ in a 2-flavor scenario ( $\nu_\mu$  and  $\nu_s$ ), MSW resonance is realized when  $V = \cos 2\theta \Delta m^2/(2E_\nu)$
- ▶ in SM, matter potential  $V$  is  $\sim 10^{-23}$  GeV; resonant oscillation length  $L_{\text{res}} \simeq \pi/(V \sin 2\theta)$  for  $\theta^2 = 10^{-3}$  reads  $L_{\text{res}} = 10^5$  km
- ▶ for KM3NeT ( $L \sim 100$  km),  $L^2/L_{\text{res}}^2 \simeq 10^{-6}$
- ▶ consider a large sterile neutrino matter potential ( $V_s \gg V_{\text{SM}}$ ) in order to have larger  $\nu_s \rightarrow \nu_\mu$  transition probability
- ▶ model with spontaneously broken  $U(1)_B$  (Pospelov, PRD 2011)

$$\mathcal{L} \supset g'_b \bar{\nu}_s \not{V} \nu_s + (g_b/3) \sum_q \bar{q} \not{V} q + \text{h.c.}$$

- ▶ EFT framework:  $\mathcal{L}_{\text{eff}} \supset \frac{g_b g'_b}{2m_V^2} [\bar{\nu}_s \gamma_\mu (1 - \gamma_5) \nu_s] [\bar{p} \gamma^\mu p + \bar{n} \gamma^\mu n]$



$$\text{Sterile neutrino potential } V_s = [g_b g'_b / m_V^2] (n_p + n_n) \equiv G_B (n_p + n_n)$$

# Neutrino production through matter-induced resonance


$$H = \begin{pmatrix} c_\theta & s_\theta \\ -s_\theta & c_\theta \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & \frac{m_s^2}{2E_\nu} \end{pmatrix} \begin{pmatrix} c_\theta & -s_\theta \\ s_\theta & c_\theta \end{pmatrix} + \begin{pmatrix} V_{\text{NC}} & 0 \\ 0 & V_s \end{pmatrix}$$

$$V_{\text{NC}} = -(\sqrt{2}/2)G_F n_n \approx -(1/2)V_{\text{CC}}$$


- ▶  $V_s = 2G_F\epsilon_{ss}(n_n + n_p)$
- ▶  $\mathcal{O}(10^2 - 10^3) \epsilon_{ss}$  considered
- ▶ for  $\mathcal{O}(100)$  PeV neutrino energy, resonance occurs for  $\sqrt{\Delta m^2} \approx m_s \simeq 2 \times 10^{-1} \sqrt{\epsilon_{ss}} \text{ keV} \Rightarrow \text{keV sterile neutrino}$




# Neutrino production through matter-induced resonance



- The difference in  $P(\nu_s \rightarrow \nu_\mu)$  implies a **larger** active neutrino flux at KM3NeT compared to that at IceCube, **alleviating the tension**




# $U(1)_B$ Constraints



► we require  $\epsilon_{ss} \simeq 100$

$$\left(\frac{g'}{g}\right)^2 \left(\frac{M_Z}{M_{Z'}}\right)^2 \simeq 100$$





Harnik, Kopp, Machado, JCAP 2012

	$U(1)_{B-L}$ (vector couplings) (Model A)	Kinetically mixed (Model B)	$U(1)_B$ (vector couplings) (Model C)
$g-2$	✓	✓	✗
Fixed Target	✓	✓	✗ <sup>a</sup>
$\Upsilon$	✓	✓	✗ <sup>b</sup>
Atomic physics	✓	✓	✗
Sun/Clusters/CAST	✓	✓	?
SN1987A	✓	✓	✓
LSW	✓	✓	✗
CMB	✓	✓	?
Borexino	✓	only if $\nu_s$ exist	✗
GEMMA	✓	✗	✗
Fifth force	✓	✗	✓

# Alleviating Sterile Neutrino Constraints

- ▶  $\nu_s$  + **secret self-interactions** (1310.6337,1806.10629)
- ▶  $\nu_s$  with initially a very large mass generated by the **VEV of a new scalar field** (1806.10629)
- ▶ Yukawa coupling of  $\nu_s$  to ultra-light scalar particle (dark matter)  
→ **large effective mass** of  $\nu_s$  in early Universe (1907.04271,1908.02278)
- ▶ **Low** reheating temperature (2501.01369)



## Model 2

# Neutrino production via non-standard interactions

- ▶ Consider the following Hamiltonian in the  $(\nu_\mu, \nu_s)$  flavor basis

$$H = \begin{pmatrix} c_\theta & s_\theta \\ -s_\theta & c_\theta \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & m_s^2/(2E_\nu) \end{pmatrix} \begin{pmatrix} c_\theta & -s_\theta \\ s_\theta & c_\theta \end{pmatrix} + \begin{pmatrix} V_{\text{NC}} & \epsilon_{\mu s} V_{\text{CC}} \\ \epsilon_{\mu s} V_{\text{CC}} & 0 \end{pmatrix}$$


- ▶ NSI term arises from  $\mathcal{L} \supset -2\sqrt{2}G_F\epsilon_{\mu s}^f (\bar{\nu}_s\gamma^\mu P_L\nu_\mu)(\bar{f}\gamma_\mu f)$
- ▶ The effective mixing angle and mass-squared difference in the limit of small  $\theta$ :

$$\theta_m = \frac{1}{2} \tan^{-1} \left( \frac{4\epsilon_{\mu s} E_\nu V_{\text{CC}}}{m_s^2 + E_\nu V_{\text{CC}}} \right) \quad \Delta m_{\text{eff}}^2 = \sqrt{(E_\nu V_{\text{CC}} + m_s^2)^2 + 16\epsilon_{\mu s}^2 E_\nu^2 V_{\text{CC}}^2}$$

- ▶  $\nu_s \rightarrow \nu_\mu$  conversion probability:

$$P_{s\mu} = \sin^2(2\theta_m) \sin^2 [\Delta m_{\text{eff}}^2 L / (4E_\nu)]$$

# Neutrino production via non-standard interactions




- ▶  $\epsilon_{\mu s}$  values up to  $\mathcal{O}(1-10)$  have a negligible effect on low-energy neutrino oscillations
- ▶ The difference in the propagation length between KM3NeT and IceCube leads to a difference in  $P_{s\mu}$ , which implies a larger active neutrino flux at KM3NeT, alleviating the tension



# Alternative explanations for the tension

- ▶ in addition to oscillations, **scattering in Earth** is an option to explain the tension



arXiv
> hep-ph > arXiv:2505.22754

High Energy Physics - Phenomenology

[Submitted on 28 May 2025]

## 'Dark' Matter Effect as a Novel Solution to the KM3-230213A Puzzle

P. S. Bhupal Dev, Bhaskar Dutta, Aparajitha Karthikeyan, Writasree Maitra, Louis E. Strigari, Ankur Verma

arXiv
> hep-ph > arXiv:2505.22711

High Energy Physics - Phenomenology

[Submitted on 28 May 2025]

## Astrophysical sources of dark particles as a solution to the KM3NeT and IceCube tension over KM3-230213A

Yasaman Farzan, Matheus Hostert

## Take-Home Message

- ▶ KM3NeT collaboration observed **highest-energy neutrino event** exceeding  $\mathcal{O}(100)$  PeV
- ▶ **No such observation has been reported at IceCube**, resulting in a tension of  $2\text{--}3.5\sigma$
- ▶ To alleviate the tension, we use the fact that the path through the Earth for KM3NeT is **an order of magnitude longer** than that for IceCube, leading to a larger  $P(\nu_s \rightarrow \nu_\mu)$  and hence a **higher flux** of muon neutrinos at KM3NeT

